Effects of Mindfulness Based Stress Reduction Therapy on Subjective Bother and Neural Connectivity in Chronic Tinnitus



Otolaryngology-Head and Neck Surgery 2015, Vol. 152(5) 919-926 © American Academy of Otolaryngology-Head and Neck Surgery Foundation 2015 Reprints and permission: sagepub.com/journalsPermissions.nav DOI: 10.1177/0194599815571556 http://otojournal.org

(\$)SAGE

Lauren T. Roland, MD¹, Eric J. Lenze, MD², Frances Mei Hardin¹, Dorina Kallogjeri, MD, MPH¹, Joyce Nicklaus¹, Andre M. Wineland, MD¹, Ginny Fendell², Jonathan E. Peelle, PhD¹, and Jay F. Piccirillo, MD¹

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

Abstract

Objective. To evaluate the impact of a Mindfulness Based Stress Reduction (MBSR) program in patients with chronic bothersome tinnitus on the (1) severity of symptoms of tinnitus and (2) functional connectivity in neural attention networks.

Study Design. Open-label interventional pilot study.

Setting. Outpatient academic medical center.

Subjects. A total of 13 adult participants with a median age of 55 years, suffering from bothersome tinnitus.

Methods. An 8-week MBSR program was conducted by a trained MBSR instructor. The primary outcome measure was the difference in patient-reported tinnitus symptoms using the Tinnitus Handicap Index (THI) and Tinnitus Functional Index (TFI) between pre-intervention, post-MBSR, and 4-week post-MBSR assessments. Secondary outcomes included change in measurements of depression, anxiety, mindfulness, and cognitive abilities. Functional connectivity magnetic resonance imaging (MRI) was performed at pre- and post-MBSR intervention time points to serve as a neuroimaging biomarker of critical cortical networks.

Results. Scores on the THI and TFI showed statistically significant and clinically meaningful improvement over the course of the study with a median Δ THI of -16 and median Δ TFI of -14.8 between baseline and 4-week follow-up scores. Except for depression, there was no significant change in any of the secondary outcome measures. Analysis of the resting state functional connectivity MRI (rs-fcMRI) data showed increased connectivity in the post-MBSR group in attention networks but not the default network.

Conclusion. Participation in an MBSR program is associated with decreased severity in tinnitus symptoms and depression

and connectivity changes in neural attention networks. MBSR is a promising treatment option for chronic bothersome tinnitus that is both noninvasive and inexpensive.

Keywords

tinnitus, functional connectivity MRI, neuroplasticity

Received October 31, 2014; revised December 23, 2014; accepted January 16, 2015.

Introduction

Subjective, idiopathic, non-pulsatile tinnitus is the perceived sensation of sound without acoustic stimulation. In the clinical setting, physicians see a striking division between bothered and nonbothered tinnitus patients that is unrelated to perceived loudness of the tinnitus.¹ Bothersome tinnitus creates problems with activities of daily living, difficulty functioning, and an inability to relax.² Tinnitus sufferers present with high rates of depression due to insomnia and problems with concentration and emotion.³ There is no reliable cure for idiopathic non-pulsatile tinnitus, and few effective treatments exist for those with bothersome tinnitus.

A major limiting factor in identifying effective therapy is a lack of understanding regarding the neurobiology of tinnitus. Although discussion of central hypotheses began long ago,⁴ most of the early work in tinnitus research focused on the cochlea. Recently, interest has shifted in a new direction

Corresponding Author:

¹Department of Otolaryngology-Head and Neck Surgery, Washington University in St Louis School of Medicine, St Louis, Missouri, USA ²Department of Psychiatry, Washington University in St Louis School of Medicine, St Louis, Missouri, USA

Lauren T. Roland, MD, Department of Otolaryngology-Head and Neck Surgery, Washington University in St. Louis School of Medicine, 660 South Euclid Ave, PO Box 8115, St. Louis, MO 63110, USA. Email: RolandL@ent.wustl.edu

to study cortical contributions to tinnitus bother.⁵ This research has been informed by resting state functional connectivity magnetic resonance imaging (rs-fcMRI),^{5,6} which indirectly measures functional neural connectivity. Recent work has used rs-fcMRI to measure connectivity differences between nonbothered tinnitus patients, bothered tinnitus patients, and healthy controls.^{7,8} Patients with bothersome tinnitus display altered patterns of neural connectivity, including key cortical attention networks, as compared to healthy controls.⁹ Nonbothered patients revealed connectivity ity networks consistent with healthy controls, suggesting that altered network connectivity may play a role in tinnitus bother.¹⁰

Mindfulness Based Stress Reduction (MBSR) is a meditation and yoga program¹¹ with a focus on mindfulness, a method for facing, exploring, and alleviating suffering by relating to present experiences. MBSR treatment reduces stress levels, improves immune function, and lowers depression and anxiety in numerous clinical settings.¹² A recent study by Gans et al¹³ provided preliminary behavioral evidence that MBSR may be an effective intervention for patients with tinnitus. Our study expands on previous work by including more participants, a follow-up test, and neuroimaging to explore the impact of MBSR on the cortical contributions of attention networks to bothersome tinnitus. We hypothesized that MBSR would improve patient symptoms by reducing level of tinnitus bother and that MBSR therapy would be associated with modulation of functional connectivity in cortical attention networks.

Methods

Design and Setting

This was an open-label, prospective, interventional trial conducted at Washington University in Saint Louis to evaluate the feasibility and effectiveness of MBSR as a treatment for patients with chronic bothersome tinnitus. A certified MBSR instructor conducted 8 weekly 2-hour MBSR sessions and one 3-hour retreat based on the MBSR workbook.¹⁴ Content included instruction in mindfulness meditation, mindful movements, and exercises to enhance mindfulness in everyday activity. Participants were given weekly homework assignments, readings, and CDs to follow during daily practice. Participants were asked to practice 30 to 45 minutes per day, and daily mindfulness logs were collected on a weekly basis. Resting state functional MRI scans were obtained at pre- and post-MBSR intervention. Questionnaire data were collected in person at pre- and post-MBSR therapy visits and electronically at 1-month following MBSR treatment using REDCap Software (Version 5.11.1). Institutional Review Board approval from the Human Research Protection Office at Washington University in Saint Louis was obtained prior to recruitment.

Participants

Eighteen adult, MBSR-naïve participants with chronic bothersome tinnitus were consented for the study between

October and November 2012. Participants were recruited through advertisements and referral from the Washington University in Saint Louis Otolaryngology clinic. Of the initial 18 participants, 2 failed screening and were excluded from the study, and 3 participants withdrew from the study for various reasons, including noncompliance with the MBSR program. The remaining 13 participants met inclusion criteria of non-pulsatile subjective unilateral or bilateral tinnitus for at least 6 months duration. In order to qualify for this study, participants needed to report global tinnitus bother as bothered more than a little but not a lot, bothered a lot, or extremely bothered when asked about the bothersome level of their tinnitus. Exclusion criteria included (1) history of head trauma, seizure, or stroke; (2) active depression or other DSM-IV axis I disorder or moderate or severe depression as evidenced by a score of 15 or higher on the Patient Health Questionnaire (PHQ-9) depression assessment¹⁵; (3) retrocochlear lesion or anatomic/structural lesion of the brain, skull base, temporal bone, or ear; (4) current use of medication for depression or anxiety; or (5) active alcohol or drug dependence.

Medical History and Behavioral Measures

Participants completed questionnaires regarding their health information, hearing history, occupational noise exposure, and tinnitus. All participants were screened for depression, anxiety, mindfulness, and cognitive abilities at the pre- and post-MBSR time points using the PHQ-9,15 Patient-Reported Outcomes Measurement Information System-Anxiety (PROMIS-Anxiety)¹⁶ evaluation, Cognitive and Affective Mindfulness Scale-Revised (CAMS-R)¹⁷ form, and Cognitive Failures Questionnaire (CFQ).18 Overall degree of tinnitus bother was assessed through the Tinnitus Global Bothersome Scale, a 5-point Likert scale ranging from a score of not bothered to extremely bothered. Selfreported measures regarding tinnitus symptoms were obtained from all participants prior to the intervention, immediately following MBSR therapy and at 4 weeks postintervention. These measures included the Tinnitus Handicap Inventory (THI)¹⁹ and the Tinnitus Functional Index (TFI).²⁰ The THI is a 25-item self-report measure with items grouped into functional, emotional, and catastrophic subscales.¹⁹ The TFI is also a 25-item questionnaire with comprehensive coverage of a wide variety of tinnitusrelated complaints. It has been shown to be highly responsive to treatment-related changes in tinnitus.²⁰

Resting State Functional Connectivity MRI

Imaging data were collected on a Siemens 3T Trio scanner. Participants were asked to relax with their eyes closed but remain awake in the scanner. Blood oxygenation level-dependent (BOLD) signal was captured using an echoplanar imaging (EPI) sequence (Repetition time [TR] = 2200 ms, echo time [TE] = 27 ms, flip angle = 90°, 4 mm isotropic voxels). Three runs of 164 volumes each were acquired for each participant. A T1-weighted structural image was also acquired (MPRAGE sequence: TR = 2100

ms, TE = 3.93 ms, flip angle = 7° , 1 × 1 × 1.25 mm voxels).

Statistical Analyses

Behavioral data were analyzed using IBM SPSS Statistics version 20.0 (IBM SPSS Inc, Chicago, Illinois, USA). The primary behavioral outcome measure was the change in tinnitus symptoms using pre- and post-MSBR differences in TFI and THI scores. Secondary behavioral outcome measures included changes in cognitive function, mindfulness, anxiety levels, and depression. As the data were not normally distributed, a Friedman test was used to assess for overall differences between the 3 paired time points. Wilcoxon signed-rank tests were conducted for post hoc analysis with the alpha level adjusted using a Bonferroni correction.

Neuroimaging Analyses

Pairs of pre- and post-MBSR intervention scans were available and usable for further data analysis for 12 subjects (an error during data preprocessing occurred for 1 participant's images). SPM8 software (Wellcome Trust Centre for Neuroimaging, London, UK) and the Automatic Analysis processing environment (version 4.1; http://github.com/rhodricusack/automaticanalysis) were used to process the MRI data.²¹ The variance between successive EPI volumes was computed, and then all EPI scans were aligned. Variance and motion parameters were used to identify scans exceeding 0.5 mm translation, 0.3° rotation, or a variance of ≥ 5 standard deviations from the mean scaled variance, which were modeled out in time-series analysis to limit the potential impact of participant motion on connectivity measures.²² EPI images for each participant were coregistered to that participant's structural image and normalized to Montreal Neurological Institute (MNI) space.²³ The data were spatially smoothed with a 9 mm full-width half maximum isotropic Gaussian kernel. Next, a time-series model was constructed that included discrete cosine basis functions to bandpass filter the data from 0.01 Hz to 0.1 Hz and the 6 movement parameters obtained from realignment, along with their squares and volterra expansions. A singular value decomposition was performed to reduce these confounds to a smaller number of regressors that explained at least 99% of the variance in the original set. Following model estimation, the residuals from the model were used as estimates of the time series of interest.

In seed-based functional connectivity analyses, the time course of brain activation is extracted from 1 region (the seed region), and the activity of all other parts of the brain are compared to this seed region to evaluate the neural connectivity. For our seed-based analyses, we chose canonical regions of interest (ROIs)²⁴ in the frontoparietal (left inferior frontal gyrus: $[-42 \ 38 \ 21]$, right inferior frontal gyrus: $[48 \ 25 \ 27]$), cingulo-opercular (left anterior frontal operculum: $[-51 \ 8 \ -2]$, right anterior frontal operculum: $[36 \ 10 \ 1]$), and default (posterior cingulate: $[8 \ -48 \ 31]$, anterior cingulate: $[12 \ 36 \ 20]$) networks. Our choice of seeds in the

frontoparietal attention network was motivated by the importance of these regions to task performance^{25,26} and implication in previous tinnitus studies,9 and our choice of seeds in the default mode network by frequent reports of MBSR-related changes in this network. Around each coordinate we extracted the mean time-series data from a sphere with a radius of 5 mm. These data were then entered in a time-series model as described earlier to identify regions of the brain in which activity was related to that of the seed region. The resulting parameter estimate maps for all participants were then entered into second-level group analyses to look at the activity for the pre-MBSR scans, post-MBSR scans, and their comparison (using a paired samples t test). For all analyses, we used a cluster-forming voxelwise threshold of P < .001, corrected for whole-brain significance at the cluster level (P < .05) using random field theory.²⁷ Results were plotted on an inflated cortical surface using Connectome Workbench (v0.85; http://www.humanconnectome.org/software/connectome-workbench.html) and slices using MRIcron (http://www.mccauslandcenter.sc.edu/ mricro/mricron/).28

Results

Participants

The 13 participants who completed the study were derived from an initial group of 70 patients who were prescreened for the study; 52 of these were excluded due to inclusion criteria or refusal to participate. Eighteen patients were consented and screened, and 2 of these patients failed screening due to medication regimens and claustrophobia. Ultimately, 16 patients enrolled in the study with 3 patients withdrawing due to personal reasons. The 13 patients who completed the study had a median age of 55 years (range, 32-65 years), and 9 (69%) of them were female. The median duration of tinnitus among the group was 4 years, with a median duration of bothersome tinnitus of 1.8 years (range, 0.6-10 years). Five patients reported that they did not have difficulty hearing speech, 7 felt that they sometimes did, and 1 reported that speech was often hard to hear. Only 1 patient reported using a hearing aid.

All 13 participants attended at least 6 MBSR sessions and 10 participants attended the retreat. Review of meditation logs indicated that participants completed an average of 35 hours of mindfulness practice during the course of the MBSR program. Three participants failed to turn in more than 2 weekly logs, so the amount of participation in mindfulness activities for these participants is unknown.

Self-reported Tinnitus Scores

TFI scores are shown in **Figure 1a**. At baseline, the median TFI score was 39 (range, 24-61), 27 at the immediate postintervention visit (range, 6-60), and 23 at the 4-week follow-up assessment (range, 4-48). At the 4-week assessment, scores were reduced as compared to baseline with a median Δ TFI of -14.8, and 8 patients (62%) had achieved a clinically meaningful reduction²⁰ of 13 points or more.



b) Tinnitus handicap inventory



Figure 1. Self-reported tinnitus health status. (A) Boxplot indicating median Tinnitus Functional Index (TFI). (B) Boxplot indicating median Tinnitus Handicap Inventory (THI). *P < .05.

Friedman's test revealed that there was a significant difference in the median Δ TFI scores, χ^2 (2 df) = 8.77, P = .012. Wilcoxon signed-rank tests revealed significant differences in TFI between pre- and post-MBSR data points (Z = -2.517, P = .012) and pre-MBSR and 4-week follow-up assessment values (Z = -2.587, P = .010). There was no significant difference between TFI values measured at the post-MBSR visit and the 4-week follow-up assessment (Z = -1.15, P = .249).

THI scores are shown in **Figure 1b**. The median THI score at baseline was 28 (range, 20-50), 28 at the immediate postintervention visit (range, 14-44), and 14 at the 4-week follow-up assessment (range, 4-32). The median change in THI score at the 4-week follow-up was -16 points from baseline. At the 4-week follow-up assessment, 10 patients (77%) had achieved a clinically meaningful difference defined by 7 points or greater reduction in THI scores.²⁹ Friedman's test revealed that there was an overall

significant difference in the Δ THI, χ^2 (2 df) = 18.12, P < .001. Individual Wilcoxon signed-rank tests, corrected for 3 samples, revealed significant differences in THI between pre- and post-MBSR values (Z = -2.635, P = .008), post-MBSR and 4-week follow-up assessment time points (Z = -2.635, P = .008), and pre-MBSR and 4-week follow-up assessment values (Z = -3.066, P = .002).

There was no significant difference in the degree of global bother assessed at both the pre- and post-MBSR visits (P = .368).

Neurocognitive Measures

Scores on the PROMIS-Anxiety, CAMS-R, and CFQ did not reveal any statistically significant change between preand postintervention time points. Although not clinically significant, the responses on the PHQ-9 assessed at the end of MBSR intervention changed from the baseline PHQ-9 scores with a median of -1 points and range from 3 points reduction to 1 point increase (P = .03). All patients had post-MBSR PHQ-9 scores of less than 15.

Functional Connectivity MRI

Seed-based functional connectivity analyses were performed on regions from 2 executive attention networks^{24,30} previously implicated in tinnitus bother^{9,10} and the default network. Functional connectivity analyses started with the left and right inferior frontal gyrus (IFG), part of the frontoparietal attention network (Figure 2). For each analysis, the seed region is displayed as a green sphere. Warm colors indicate brain regions that show significant connectivity with the seed region, with the colormap corresponding to the t statistic. Connectivity with the left IFG was significantly increased following MBSR in 3 clusters: left superior temporal gyrus (peak [-58 - 6 - 2], peak Z = 4.41, 518 voxels), posterior cingulate (peak [-12 - 50 20], peak Z = 4.15, 194 voxels), and visual cortex (peak [8 - 84 - 14], peak Z = 3.83, 179 voxels). Connectivity was increased with the right IFG to a cluster in the posterior cingulate cortex (peak [-12 - 50 22], peak Z = 4.14, 283 voxels). There were no regions showing significant decreases in connectivity.

Similar analyses were conducted using seed regions from the left and right anterior operculum (aO), part of the cingulo-opercular network associated with task set maintenance (**Figure 3**). Significant increases from left aO were found to a cluster extending into the left cingulate cortex (peak [-16 - 640], peak Z = 4.15, 253 voxels); increases from right aO were found to a cluster spanning left temporal and parietal cortex (peak [-36 - 4222], peak Z = 5.00, 432voxels), right frontal regions bordering anterior cingulate and middle frontal gyrus (peak [202024], peak Z = 4.25, 192 voxels), and left superior temporal gyrus (peak [-54-2-4], peak Z = 3.85, 151 voxels). There were no regions showing significant decreases in connectivity.

Lastly, the default network was investigated using seeds in posterior cingulate cortex (PCC) and anterior cingulate cortex (ACC) (**Figure 4**). Although connectivity with both



Figure 2. Functional connectivity for inferior frontal gyrus regions of interest pre and post Mindfulness Based Stress Reducation (MBSR). The right column shows post > pre functional connectivity data with significant clusters circled in white.



Figure 3. Functional connectivity for left and right anterior operculum regions of interest pre and post Mindfulness Based Stress Reduction (MBSR). The right panel illustrates significant clusters showing increased connectivity for post-MBSR relative to pre-MBSR.

seeds revealed a stereotypical pattern of default network connectivity, there were no statistically significant differences between pre- and post-MBSR connectivity with these seeds.

Discussion

This study examined changes in symptoms and neural connectivity after MBSR in patients with bothersome tinnitus. A significant and clinically meaningful reduction in tinnitus bother was observed following MBSR using both THI and TFI assessment measures. Measures of depression, anxiety, mindfulness, and cognitive abilities did not show clinically meaningful changes after MBSR, although there was a statistically significant change in pre-post depression scores. A seedbased analysis performed on rs-fcMRI scans obtained before and after MBSR treatment revealed changes in functional connectivity within regions associated with known attention networks. Together, these results suggest that the subjective



Figure 4. Functional connectivity for posterior and anterior cingulate regions of interest pre and post Mindfulness Based Stress Reduction (MBSR). There were no significant differences between pre- and post-MBSR scans.

improvement associated with participation in an MBSR program may be related to altered neural connectivity.

Previous behavioral studies done at other institutions have reported preliminary evidence for the efficacy of MBSR in reducing bother associated with chronic tinnitus.^{13,31} Our current findings expand on previous studies through the use of a larger sample size and the incorporation of advanced neuroimaging to study the functional connectivity within cortical attention networks. The interpretation of prior studies investigating the effects of MBSR using rs-fcMRI is limited by variability in methodology and analysis approach.

Our findings agree with past studies, which are largely consistent in reporting changes in neural connectivity following MBSR. However, studies using rs-fcMRI to investigate the effects of MBSR show considerable heterogeneity in the populations studied, seed regions, and analysis approach, making direct comparisons of specific neural changes difficult. Several studies have focused exclusively on the default mode network.32-34 Hasenkamp and Barsalou³⁵ found that participants with more meditation experience exhibited increased connectivity within attention networks as well as between attentional regions and the default mode network. In another study examining the use of MBSR in healthy adults, differences in functional connectivity between the MBSR and control groups were found in the auditory/salience and medial visual networks, but there were no noted changes in the default mode network or attention networks.³⁶ Thus, although there is agreement that MBSR results in changed neural connectivity, the consistency of MBSR effects on specific attention networks across populations from prior studies is difficult to evaluate.

The functional connectivity results of this study suggest multiple changes and increased activation between networks

responsible for attention and focus. Our previous work⁹ suggests that patients with bothersome tinnitus have abnormal functional connectivity in networks responsible for attention, decision making, and somatosensory input. The functional connectivity observations in the current study suggest that participation in an MBSR program can result in neuroplastic changes in attention networks. These changes coincided with self-reported improvement in symptoms.

There are several limitations to this study. First, this was a pilot study and we did not use a control group. An appropriate control group would be matched in regards to age, gender, and tinnitus bother and would undergo an intervention that does not incorporate the techniques of MBSR but does control for the attention provided to the participant during MBSR. A health education program (HEP)³⁷ has been suggested as an appropriate active control condition for MBSR: while HEP improves chronic disease management, it does not teach mindfulness techniques or address worry and rumination and is thus a credible behavioral intervention that has none of the posited active ingredients of MBSR.38 A second limitation of our study was the small sample size (n = 13). The participants represented a wide range of ages, variability in pre- and post-MBSR tinnitus symptom scores, amount of time spent using the MBSR program at home, and attendance at weekly sessions. Due to our limited sample size, we were not able to adequately control for these factors or conduct cogent subgroup analyses. Future studies including both an active control group and a larger sample size will be beneficial.

Conclusion

This research identified a reduction in tinnitus symptoms and neuroimaging changes in functional connectivity in cortical attention networks, coinciding with clinical improvement of symptoms. MBSR appears to be a beneficial technique for tinnitus patients: MBSR is inexpensive, noninvasive, and takes advantage of the neuroplasticity of the brain to potentially create lasting benefits.

Acknowledgments

The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Author Contributions

Lauren T. Roland, analysis and interpretation of data, drafting of work, final approval, agreement to be accountable; Eric J. Lenze, study concept, critical revision, final approval, agreement to be accountable; Frances Mei Hardin, data acquisition, drafting of work, final approval, agreement to be accountable; Dorina Kallogjeri, study concept, analysis and interpretation of data, drafting of work, final approval, agreement to be accountable; Joyce Nicklaus, study design, data acquisition, critical revision, final approval, agreement to be accountable; Andre M. Wineland, study concept, critical revision, final approval, agreement to be accountable; Ginny Fendell, study concept, data acquisition, final approval, agreement to be accountable; Jonathan E. Peelle, analysis and interpretation of data, drafting of work, final approval, agreement to be accountable; Jay F. Piccirillo, study concept/design, interpretation of data, drafting of work, final approval, agreement to be accountable.

Disclosures

Competing interests: Eric J. Lenze, Roche and Lundbeck grant funding; Ginny Fendell, independent consultant, teaches Mindfulness Based Stress Reduction course.

Sponsorships: None.

Funding source: This research and publication was supported, in part, by the Washington University Institute of Clinical and Translational Sciences grant UL1 TR000448 from the National Center for Advancing Translational Sciences, the "Development of Clinician/Researchers in Academic ENT" T32 DC00022 from the National Institutes of Deafness and Other Communication Disorders, and grant AT007064 from the National Center for Complementary and Alternative Medicine. This project was also supported, in part, with funding from the Mallinckrodt Institute of Radiology at the Washington University School of Medicine. The use of REDCap software was supported by Clinical and Translational Science Award (CTSA) Grant UL1TR00048 and Siteman Comprehensive Cancer Center and NCI Cancer Center Support Grant P30 CA091842.

References

- 1. Coles RRA, Hallam RS. Tinnitus and its management. *Br Med Bull*. 1987;43:983-998.
- Hoffman HJ, Reed GW. Epidemiology of tinnitus. In: Snow JB Jr, ed. *Tinnitus: Theory and Management*. Hamilton, Ontario: BC Decker, Inc; 2004: 16-41.
- Dobie R. Overview: suffering from tinnitus. In: Snow JB Jr, ed. *Tinnitus: Theory and Management*. Hamilton, Ontario: BC Decker; 2004: 1-7.
- 4. Tyler RS, Conrad-Armes D. Masking of tinnitus compared to masking of pure tones. *J Speech Hear Res.* 1984;27:106-111.

- Husain FT, Schmidt SA. Using resting state functional connectivity to unravel networks of tinnitus. *Hear Res.* 2014; 307:153-162.
- Buckner RL, Krienen FM, Yeo BTT. Opportunities and limitations of intrinsic functional connectivity MRI. *Nat Neurosci*. 2013;16:832-837.
- Maudoux A, Lefebvre P, Cabay JE, et al. Auditory restingstate network connectivity in tinnitus: a functional MRI study. *PLoS One*. 2012;7:e36222.
- 8. Maudoux A, Lefebvre P, Cabay JE, et al. Connectivity graph analysis of the auditory resting state network in tinnitus. *Brain Res.* 2012:1485;10-21.
- Burton H, Wineland A, Bhattacharya M, Nicklaus J, Garcia K, Piccirillo J. Altered networks in bothersome tinnitus: a functional connectivity study. *BMC Neurosci.* 2012;13:1-15.
- Wineland AM, Burton H, Piccirillo J. Functional connectivity networks in nonbothersome tinnitus. *Otolaryngol Head Neck Surg.* 2012;147:900-906.
- 11. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol*. 2003;10:144-156.
- Lengacher CA, Johnson-Mallard V, Post-White J, et al. Randomized controlled trial of Mindfulness-Based Stress Reduction (MBSR) for survivors of breast cancer. *Psycho-Oncology*. 2009;18:1261-1272.
- Gans J, O'Sullivan P, Bircheff V. Mindfulness based tinnitus stress reduction pilot study: a symptom perception shift program. *Mindfulness*. 2013;5:1-12.
- Stahl B, Goldstein E. Mindfulness-Based Stress Reduction Workbook. Oakland, CA: New Harbinger Publications, Inc; 2010.
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. J Gen Intern Med. 2001; 16:606-613.
- Reeves DL, Winter KP, Bleiberg J, Kane RL. ANAM genogram: historical perspectives, description, and current endeavors. *Arch Clin Neuropsychol.* 2007;22(suppl 1):S15-S37.
- Feldman G. Mindfulness and emotion regulation: the development and initial validation of the Cognitive and Affective Mindfulness Scale-Revised (CAMS-R). *Psychopathol Behav* Assess. 2007;29:177-190.
- Broadbent DE, Cooper PF, Fitzgerald P, Parkes KR. The Cognitive Failures Questionnaire (CFQ) and its correlates. *Br J Clin Psychol.* 1982;21:1-16.
- Newman CW, Jacobson GP, Spitzer JB. Development of the Tinnitus Handicap Inventory. Arch Otolaryngol Head Neck Surg. 1996;122:143-148.
- Meikle MB, Henry JA, Griest SE, et al. The Tinnitus Functional Index: development of a new clinical measure for chronic, intrusive tinnitus. *Ear Hear*. 2012;33:153-176.
- 21. Cusak R, Vincente-Grabovetsky A, Mitchell DJ, et al. Automatic analysis (aa): Efficient neuroimaging workflows and parallel processing using Matlab and XML. *Front Neuroinform*. 2015;8:90.
- Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*. 2012;59:2142-2154.

- Ashburner J, Friston KJ. Unified segmentation. *NeuroImage*. 2005;26:839-851.
- 24. Power JD, Cohen AL, Nelson SM, et al. Functional network organization of the human brain. *Neuron*. 2011;72:665-678.
- 25. Power JD, Petersen SE. Control-related systems in the human brain. *Curr Opin Neurobiol.* 2013;23:223-228.
- Duncan J. The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends Cogn Sci.* 2010;14:172-179.
- Worsley KJ, Evans AC, Marrett S, Neelin P. A threedimensional statistical analysis for CBF activation studies in human brain. J Cereb Blood Flow Metab. 1992;12:900-918.
- Rorden C, Brett M. Stereotaxic display of brain lesions. *Behav Neurol.* 2000;12:191-200.
- Zemen F, Koller M, Figueiredo R, et al. Tinnitus Handicap Inventory for evaluating treatment effects: which changes are clinically relevant? *Otolaryngol Head Neck Surg.* 2011;145: 282-287.
- Petersen SE, Posner MI. The attention system of the human brain: 20 years after. *Annu Rev Neurosci*. 2012;35:73-89.
- Kreuzer PM, Goetz M, Holl M, et al. Mindfulness-and bodypsychotherapy-based group treatment of chronic tinnitus: a randomized controlled pilot study. *BMC Complement Altern Med.* 2012;12:235.

- Taylor VA, Daneault V, Grant J, et al. Impact of meditation training on the default mode network during a restful state. Soc Cogn Affect Neurosi. 2012;8:4-14.
- Brewer JA, Worhunsky PD, Gray JR, Tang YY, Weber J, Kober H. Meditation experience is associated with differences in default mode network activity and connectivity. *Proc Natl Acad Sci USA*. 2011;108:20254-20259.
- Wells RE, Yeh GY, Kerr CE, et al. Meditation's impact on default mode network and hippocampus in mild cognitive impairment: a pilot study. *Neurosci Lett.* 2013;556:15-19.
- Haskenkamp W, Barsalou LW. Effects of meditation experience on functional connectivity of distributed brain networks. *Front Hum Neurosci.* 2012;6:38.
- Kilpatrick LA, Suyenobu BY, Smith SR, et al. Impact of Mindfulness Based Stress Reduction training on intrinsic brain connectivity. *Neuroimage*. 2011;56:290-298.
- Lorig K, Holman H, Sobel D, Laurent D, Gonzalez V. Living a Healthy Life with Chronic Conditions: For Ongoing Physical and Mental Health Conditions. Boulder, CO: Bull Publishing Company; 2007.
- MacCoon DG, Imel ZE, Rosenkranz MA, et al. The validation of an active control intervention for Mindfulness Based Stress Reduction (MBSR). *Behav Res Ther.* 2012;50:3-12.